

Heavy probes of initial conditions & novel observables

Sandeep Chatterjee
IISER Berhampur

Hard Probes in non-equilibrium QCD matter,
16-27 March, 2026
ICTS Bengaluru


Heavy probes of initial **geometry, EM fields** & **novel observables**

Sandeep Chatterjee
IISER Berhampur


Hard Probes in non-equilibrium QCD matter,
16-27 March, 2026
ICTS Bengaluru

Initial Collision Geometry




Peripheral Collision



(near) Central Collision




Initial Collision Geometry - elliptic flow





Initial Collision Geometry - directed flow

Initial Collision Geometry - directed flow



+



Initial Collision Geometry - directed flow



d+Au at $\sqrt{s_{NN}} = 200$ GeV (PHOBOS 2011)


FB asymmetric deposition by a participant source

Initial Collision Geometry - directed flow




d+Au at $\sqrt{s_{NN}} = 200$ GeV (PHOBOS 2011)

FB asymmetric deposition by a participant source




Tilted fireball !!

Initial Collision Geometry - directed flow




Tilted fireball !!




Au+Au at $\sqrt{s_{NN}} = 200$ GeV (STAR 2009)

Initial Collision Geometry - directed flow

(a) reaction plane



Tilted fireball !!




Bozek, Wyskiel 2010


Heavy probe of Initial Tilt

HQ from hard processes \rightarrow FB-symmetric

Rapidity-even HQ dragged by Rapidity-odd bulk



Bulk vs heavy flavor



charm, anti-charm stronger probes of the tilt than the light flavor




Heavy probe of Initial Tilt

need to calibrate



- the tilt of the bulk: constrained by charged particle v_1 , Božek, Wyskiel 2010
- drag between the bulk and heavy flavor: constrained by heavy flavor R_{AA} and v_2 at mid-rapidity, we use an ansatz

$$\gamma = \gamma_0 T \left(\frac{T}{m} \right)^x$$



Heavy probe of Initial Tilt - Calibrating the drag




Heavy probe of initial tilt: 10 times larger v_1 than light flavor!!




Heavy probe of initial tilt: 10 times larger v_1 than light flavor!!



Heavy probe of initial tilt: 10 times larger v_1 than light flavor!!




SC, Bozek 2018



**Tilt ~ Angular Momentum -> Heavy probe of the Initial Angular momentum:
Refer to Amaresh's Talk**


Initial Electromagnetic field



- $\mathbf{B} = -B \hat{y}$, B decreases as the spectators recede. For non-zero conductivity σ of the medium, this gives rise to a clock-wise \mathbf{E} in the above reaction plane
- A positive charge at $\eta > 0$ experiences \mathbf{B} force along \hat{x} while \mathbf{E} force along $-\hat{x}$, the net force resulting in a directed flow v_1

Light probe of Initial Electromagnetic field


v_1 split between positive and negative charged particles due to EM field



Gursoy, Kharzeev, Rajagopal 2014



Heavy probe of Initial Electromagnetic field

1000 times stronger effect on heavy flavor !!



Das, Plumari, SC, Alam, Scardina, Greco 2016

Heavy probe of initial Electromagnetic field



Alice measurements hints at disagreement

Heavy probes of initial geometry, EM fields & novel observables

With Kaiser Shafi, Tribhuban Parida, *in preparation*

Motivation

Recall this previous slide: T dep. of γ is not well understood

need to calibrate

- the tilt of the bulk: constrained by charged particle v_1 , Božek, Wyskiel 2010
- drag between the bulk and heavy flavor: constrained by heavy flavor R_{AA} and v_2 at mid-rapidity, we use an ansatz

$$\gamma = \gamma_0 T \left(\frac{T}{m}\right)^x \quad \mathbf{x \text{ is not well constrained}}$$

Phase space of hadrons in the final state

- In the lab frame: (p_x, p_y, p_z)
- For most observables,
region of interest:
(transverse spectra,
elliptic flow) $\text{mid-rapidity, i.e. } p_z \sim 0$
- Resulting 2-D phase space: $(p_x, p_y) \rightarrow (p_T, \phi)$



thus, we talk of 'p_T' dependence and 'azimuthal dependence'

Azimuthal dependence

Periodicity in ϕ provides a natural basis
(Fourier expansion):

$$N(\phi) = N_0 + v_1 \cos(\phi) + \dots \text{ with } v_n \ll 1;$$


p_T dependence



- p_T dependence by eye comparison!
Are we missing important physics here?

Motivates us to revisit the p_T dependence

2-D mode decomposition: Fourier-Bessel Basis



ρ_{τ} basis functions

Bessel functions: eigenfunctions of Laplacian on disc \rightarrow provides a natural basis for the ρ_{τ} dependence

$$\phi_{m,n}(r, \theta) := \frac{1}{J_{|m|+1}(\lambda_{m,n})} J_m \left(\frac{r}{r_0} \lambda_{m,n} \right) e^{im\theta}$$

J \rightarrow Bessel function of first kind

$$f(r, \theta) = \sum_{m,n} A_{m,n} \phi_{m,n}(r, \theta)$$


- 2-D mode expansion

- *J. Phys. G*, vol. 40, 2013, p. 095 103


Drag coefficient

$$\gamma = \gamma_0 T (T/m)^x$$


γ vs T



Pb+Pb 5.02 TeV, D_0 : Glauber+MUSIC+Langevin, 20-30%




p_T mode decomposition of v_2



T dependence is revealed !!

p_T mode decomposition of v_2




T dependence is revealed !!

Summarising..


Heavy flavor offers unique sensitive probes to understand the initial

- 3 dimensional distribution of the fireball
- Electromagnetic fields



Summarising..

Introduced a set of novel observables that provides a complete **2-D decomposition of the transverse phase space distribution** - demonstrated their utility in **extracting the temperature dependence of the heavy quark drag coefficient**



Thank You