



SOme illustrative results for Lecture 3

Based on 2506.19194, 2402.04488



A model potential in 1D

Need to design a realistic 1D bona fide potential $V + iW$ (based on 3D IQCD results, tuned to reproduce 3D mass spectra and decay widths)



1D potential from R. Katz, S. Delorme & PBG, Eur. Phys. J. A (2022) 58:198

Typical regimes in OQS



Some tests of semi-classical approximations
(abelian case)


Based on 2506.19194

Typical evolution of the density matrix

Reduced density matrix of a $Q\bar{Q}$ pair: $\langle x_1, x_2 | \mathcal{D} | x'_1, x'_2 \rangle = \mathcal{D}(x_1, x_2; x'_1, x'_2)$

$$\mathcal{D}(x_1, x_2; x'_1, x'_2) \mapsto \mathcal{D}(s, s')$$


$$s = x_1 - x_2 \quad s' = x'_1 - x'_2$$




Initial wave packet

$$\psi(x) = \left(\frac{1}{\pi\sigma^2} \right)^{1/4} e^{-\frac{x^2}{2\sigma^2}} \quad (\sigma = 0.38 \text{ fm})$$

Wigner distribution
(‘phase space’ distribution)




Evolution in QGP with $T=300$ MeV




Looking at finer details

$$d(t) = \int \frac{dr dp}{2\pi\hbar} |\mathcal{D}_{\text{QM}}(t, r, p) - \mathcal{D}_{\text{SC}}(t, r, p)|$$



Impact of the non-unitary dynamics




SOme results for non abelian case

Based on 2402.04488

Collisional decoherence


charm quark-antiquark pair at $T=300$ MeV



[From S.Delorme, R. Katz, T. Gousset, PB. Gossiaux, JPB (2023)]

Color equilibration

$$\rho_{s,o} = \langle r | \mathcal{D}_{s,o} | r \rangle$$



[From S. Delorme, R. Katz, T. Gousset, PB. Gossiaux, JPB (2023)]

Asymptotic state

