

# Direct Photon Production from Concurrent Minijet-Hydrodynamics Evolution

Soham Banerjee   Mayank Singh   Daniel Pablos Alfonso   Jean-Francois Paquet  
Sangyong Jeon   Charles Gale

March 23, 2026



## Motivation and Overview

- Jets with relatively low- $p_T$  (mini-jets)  $\rightarrow$  can continuously deposit energy-momentum into the medium
- Direct photons are probes of the quark-gluon plasma (QGP)
- They are emitted throughout the space-time evolution and escape largely unmodified
- How does continuous minijet(hard) energy loss affect direct photon observables?
- We investigate how a concurrent minijet+hydro evolution impacts:
  - Direct photon spectra
  - Photon elliptic flow ( $v_2$ )
- Framework: IP-Glasma + MUSIC + UrQMD + dedicated photon module, with minijet dynamics from a hybrid model

## Introduction to Minijets:

- Hard parton scattering  $\rightarrow$  outgoing partons  $\rightarrow$  shower/fragment  $\rightarrow$  jets<sup>1,2</sup>
- Mini-jets with relatively low transverse momentum:

$$p_T \sim \mathcal{O}(4-10) \text{ GeV}$$

- Produced in initial hard pQCD scatterings
- Intermediate regime:
  - between soft bulk medium and rare high- $p_T$  jets
- If  $x_{\text{stop}} > \tau_0$ :

$\Rightarrow$  slow thermalization  $\Rightarrow$  persistent energy-momentum source during hydro



Wake, Credit: Edmont, Wikimedia Commons, CC BY-SA 3.0

<sup>1</sup> Busza, Rajagopal, van der Schee, *Ann.Rev.Nucl.* (2018)

<sup>2</sup> Solana, Gulhan, Milhano, Pablos, Rajagopal, *JHEP* (2014)

## Choice of the jet transverse momentum




Figure: Ref. Pablos, Singh, Jeon, Gale, PRC (2022)

# Modified Hydro Evolution




Figure: Snapshots of the 3D isotherms. Ref. Pablos, Singh, Jeon, Gale, PRC (2022)

# Minijets: How They Affect the Medium

- Continuous energy–momentum deposition into the QGP fluid
- Induces *spikier* hydrodynamic fields:
  - Additional event-by-event fluctuations
- Affects final-state multiplicity:
  - Mini-jets add entropy
  - To keep realistic  $dN_{ch}/d\eta$ , initial entropy must be reduced
- Minijet momenta are randomly oriented:
  - $\Rightarrow$  tend to reduce collective flow
- Impact on transport extraction:
  - To reproduce measured  $v_n$ ,
  - Presence of minijets favors lower QGP  $\eta/s$

| $p_J^{\min}$ [GeV] | $s_{\text{factor}}$ | $\eta/s$ |
|--------------------|---------------------|----------|
| 4                  | 0.45                | 0.02     |
| 7                  | 0.82                | 0.10     |
| 10                 | 0.90                | 0.125    |
| No jets            | 0.915               | 0.13     |

- Mini-jets require adjusting:
  - initial entropy
  - shear viscosity
- Direct photons can help constrain these effects

<sup>1</sup> Pablos, Singh, Jeon, Gale, PRC (2022)


# Direct Photons as Probes of the QGP

- Categorized according to their origins <sup>1</sup>:
  - Prompt Photons  $\implies$  Produced in initial hard partonic scatterings (e.g., Compton scattering  $qg \rightarrow q\gamma$  and annihilation  $q\bar{q} \rightarrow g\gamma$ )
  - Thermal Photons  $\implies$  QGP  $\oplus$  hadronic gas
  - Decay Photons  $\implies$  hadron decays (subtracted experimentally)
- Direct photons are defined as:

$$\text{Direct photons} = \text{Prompt photons} \oplus \text{Thermal photons}$$

<sup>1</sup> Jean-François Paquet, Thesis (2016)

# Simulation Framework



# Hybrid Strong/Weak Coupling

- Scale separation <sup>1</sup>
  - Hard jet production and splittings: perturbative QCD (DGLAP)
  - Jet-medium interactions: strongly coupled dynamics
- Model assumptions
  - Soft exchanges
  - Vacuum-like splittings: DGLAP splitting probabilities

## Energy-loss rate <sup>2</sup>

$$\frac{1}{E_{\text{in}}} \frac{dE}{dx} = - \frac{4x^2}{\pi x_{\text{stop}}^2 \sqrt{x_{\text{stop}}^2 - x^2}}$$

## Stopping distance

$$x_{\text{stop}} = \frac{1}{2 \kappa_{\text{sc}}} \frac{E_{\text{in}}^{1/3}}{T^{4/3}}$$

<sup>1</sup> Chesler and Rajagopal, PRD (2014)

<sup>2</sup> Solana, Gulhan, Milhano, Pablos, Rajagopal, JHEP(2014)

# IP-Glasma Initial Conditions with Concurrent Minijets

- **Bulk initial state from IP-Glasma<sup>1,2</sup>:**

- CGC-based description of the colliding nuclei
- Generates fluctuating energy–momentum density profiles
- Hydrodynamic initialization at

$$\tau_0 = 0.4 \text{ fm}$$

- **Mini-jets from hard scatterings:**

- Produced by PYTHIA

- **Concurrent minijet–hydrodynamic evolution:**

- Hybrid strong/weak-coupling description of jet propagation
- Lost energy is deposited into the medium as a source term<sup>3</sup>:

$$\partial_\mu T_{\text{fluid}}^{\mu\nu} = J_{\text{minijets}}^\nu$$

<sup>1</sup> Scott McDonald, Thesis (2020)

<sup>2</sup> Gelis, Iancu, Jalilian-Marian, Venugopalan, ARNPS (2010)

<sup>3</sup> Pablos, Singh, Jeon, Gale, PRC (2022)

# How We Calculate Direct Photons

- **Prompt photons**

- Computed using a baseline prompt photon spectrum from  $p+p$  collisions
- Scaled to Pb+Pb collisions using binary collision scaling:

$$\left. \frac{dN_{\text{prompt}}^{\gamma}}{dp_{\tau}} \right|_{AA} = \left. \frac{dN_{\text{prompt}}^{\gamma}}{dp_{\tau}} \right|_{pp} \times N_{\text{coll}}$$

- **Thermal photons**


- The full hydrodynamic evolution is stored:  $T(x)$ ,  $u^{\mu}(x)$ ,  $\pi^{\mu\nu}(x)$ , etc
- Local thermal emission rates are evaluated at each space-time point
- The total thermal yield is obtained by integrating the rates over the 4-volume:

$$E \frac{dN_{\text{th}}^{\gamma}}{d^3p} = \int d^4x \left( E^* \frac{dR^{\gamma}}{d^3p^*} \right) (T(x), u^{\mu}(x))$$


- Emission is terminated at a temperature:

$$T_{\text{sw}} = 105 \text{ MeV}$$

# Preliminary Results: Photon Spectra (Central)




# Preliminary Results: Photon Elliptic Flow $v_2$



# Photon $v_2$ for (0-40)% w/w.o $\delta f$ correction



## Thermal photon $v_2$ for 0-40% cent





# Summary

- We studied direct photon production in a hybrid minijet+hydro framework:
  - IP-Glasma + MUSIC + UrQMD
  - Minijets from PYTHIA with hybrid strong/weak-coupling energy loss
- Mini-jets significantly impact:
  - Hydrodynamic fields and fluctuations
  - Photon yields and elliptic flow
  - Interpretation of QGP transport coefficients
- Direct photons provide an effective probe of minijet dynamics and medium evolution



# Comparison with previous results




# Comparison with previous results 1:




## Comparison with previous results 2:



# Preliminary Results: Photon Spectra (Semi-central)



# Evolution of energy density-weighted average temperature



# Evolution of energy density-weighted average momentum anisotropy

