

Primordial Black Holes from the Supercooled Phase Transitions with Radiative Symmetry Breaking

Francesco Rescigno

University of Rome Tor Vergata, INFN Roma2

arXiv:2412.06889 [hep-ph] I.K. Banerjee, F.Rescigno, A.Salvio

Hearing beyond the standard model with cosmic sources of Gravitational Waves
ICTS-TIFR, Bengaluru, January 2025



- 1 FOPT, Supercool Expansion and Improved Supercool Expansion
- 2 Late blooming mechanism

1 FOPT, Supercool Expansion and Improved Supercool Expansion

2 Late blooming mechanism

First Order Phase Transitions (FOPT)

- FOPT can occur in the early Universe when the temperature T drops below a critical value T_c
- The finite temperature effective potential $V(\chi, T)$ develops a new stable minima (**true vacuum**), while the old minima becomes metastable (**false vacuum**)
- The false vacuum eventually **decay in the true vacuum**
- The false vacuum decay manifest as the **nucleation of true vacuum bubbles** in a background of false vacuum




Figure 1: Image from JCAP **04** (2023), 051 [arXiv:2302.10212 [hep-ph]], A. Salvio

Radiative Symmetry Breaking (RSB)

- We start from the most general no-scale Lagrangian:

$$\mathcal{L}_{\text{ns}} = -\frac{1}{4}F_{\mu\nu}^A F_A^{\mu\nu} + \frac{1}{2}D_\mu\phi_a D^\mu\phi_a + \bar{\psi}_j i\not{D}\psi_j - \frac{1}{2}(Y_{ij}^a \psi_i \psi_j \phi_a + \text{h.c.}) - V_{\text{ns}}(\phi),$$

$$D_\mu\phi_a = \partial_\mu\phi_a + i\theta_{ab}^A V_\mu^A \phi_b, \quad D_\mu\psi_j = \partial_\mu\psi_j + it_{jk}^A V_\mu^A \psi_k,$$

$$V_{\text{ns}}(\phi) = \frac{\lambda_{abcd}}{4!} \phi_a \phi_b \phi_c \phi_d.$$

Radiative Symmetry Breaking (RSB)

- At quantum level the couplings depend on the RG energy μ , there may be some specific value $\mu = \tilde{\mu}$ at which the potential $V_{\text{ns}}(\phi)$ develops a flat direction parametrized as

$$\phi_a = \chi \nu_a$$

$$V(\chi) = \frac{\lambda_\chi(\mu)}{4} \chi^4, \quad \lambda_\chi(\mu) \equiv \frac{1}{3!} \lambda_{abcd}(\mu) \nu_a \nu_b \nu_c \nu_d,$$

such that $\lambda_\chi(\tilde{\mu}) = 0$ (flat direction).

- The one-loop effective potential at zero temperature is

$$V_q(\chi) = \frac{\lambda_\chi(\mu)}{4} \chi^4 + \frac{\beta_{\lambda_\chi}}{4} \left(\log \frac{\chi}{\mu} - \frac{1}{4} \right) \chi^4, \quad \beta_{\lambda_\chi} \equiv \mu \frac{d\lambda_\chi}{d\mu}.$$

Radiative Symmetry Breaking (RSB)

- Setting $\mu = \tilde{\mu}$ we obtain $\lambda_\chi = 0$, i.e.

$$V_q(\chi) = \frac{\bar{\beta}}{4} \left(\log \frac{\chi}{\chi_0} - \frac{1}{4} \right) \chi^4,$$

$$\bar{\beta} \equiv [\beta_{\lambda_\chi}]_{\mu=\tilde{\mu}}, \quad \chi_0 \equiv \frac{\tilde{\mu}}{e^{1/4+a_s}}.$$

- When the conditions,

$$\begin{cases} \lambda_\chi(\tilde{\mu}) = 0 & \text{(flat direction)} \\ \beta_{\lambda_\chi}(\tilde{\mu}) > 0 & \text{(minimum condition)} \end{cases}$$

are fulfilled χ_0 is the zero temperature vacuum-expectation value of χ and is the new absolute minima of $V(\chi)$

- the fluctuation around χ_0 have mass $m_\chi^2 = \bar{\beta}\chi_0^2$

Radiative Symmetry Breaking (RSB)

- The non trivial minimum can generically break global and/or local symmetries and thus generate the particle masses, where χ_0 plays the role of symmetry breaking scale

$$\mathcal{L}_{\chi h} \equiv \frac{1}{2} \lambda_{ab} \phi_a \phi_b |\mathcal{H}|^2, \quad \lambda_{\chi h}(\mu) \equiv \lambda_{ab}(\mu) \nu_a \nu_b.$$

- RG improving and setting $\mu = \tilde{\mu}$

$$\mathcal{L}_{\chi \mathcal{H}} = \frac{1}{2} \lambda_{\chi \mathcal{H}}(\tilde{\mu}) \chi^2 |\mathcal{H}|^2,$$

evaluated at the minimum

$$\mu_h^2 = \frac{1}{2} \lambda_{\chi h}(\tilde{\mu}) \chi_0^2.$$

Effective thermal potential

- For the no-scale theory in question, we have a thermal effective potential of the form:

Effective Thermal Potential

$$V_{\text{eff}}(\chi, T) = V_q(\chi) + \frac{T^4}{2\pi^2} \left(\sum_b n_b J_B(m_b^2(\chi)/T^2) - 2 \sum_f J_F(m_f^2(\chi)/T^2) \right) + \Lambda_0,$$

$$J_B(x) \equiv \int_0^\infty dp p^2 \log \left(1 - e^{-\sqrt{p^2+x}} \right) = -\frac{\pi^4}{45} + \frac{\pi^2}{12}x - \frac{\pi}{6}x^{3/2} - \frac{x^2}{32} \log \left(\frac{x}{a_B} \right) + O(x^3),$$

$$J_F(x) \equiv \int_0^\infty dp p^2 \log \left(1 + e^{-\sqrt{p^2+x}} \right) = \frac{7\pi^4}{360} - \frac{\pi^2}{24}x - \frac{x^2}{32} \log \left(\frac{x}{a_F} \right) + O(x^3),$$

$$a_B = 16\pi^2 \exp(3/2 - 2\gamma_E), \quad a_F = \pi^2 \exp(3/2 - 2\gamma_E).$$

- In such models the production of true vacuum bubbles is described by the **time independent bounce**

Decay rate of the false vacuum per unit of volume (time independent bounce)

$$\Gamma \approx T^4 \exp(-S_3/T),$$

$$S_3 \equiv 4\pi \int_0^\infty dr r^2 \left(\frac{1}{2} \chi'^2 + \bar{V}_{\text{eff}}(\chi, T) \right) = -8\pi \int_0^\infty dr r^2 \bar{V}_{\text{eff}}(\chi, T),$$

$$\chi'' + \frac{2}{r} \chi' = \frac{d\bar{V}_{\text{eff}}}{d\chi}, \quad \chi'(0) = 0, \quad \lim_{r \rightarrow \infty} \chi(r) = 0, \quad \bar{V}_{\text{eff}}(\chi, T) = V_{\text{eff}}(\chi, T) - V_{\text{eff}}(0, T).$$

Properties of Phase Transition in RSB models

- Phase Transitions in RSB models are **always FOPT**
- FOPTs in RSB feature always a **supercooling phase**, i.e. the temperature drops much below T_c before the bubble nucleation became effective. This ensures that:
 - ▶ at relevant temperatures the **thermal corrections are small**
 - ▶ the zero temperature mass is much larger than the thermal mass, **avoiding the infrared Linde problem**
- FOPT in RSB models are always **strong**, the PT release a large amount of energy




Figure 2: Image from JCAP **04** (2023), 051 [arXiv:2302.10212 [hep-ph]], A. Salvio

JCAP **04** (2023), 051 [arXiv:2302.10212 [hep-ph]], A. Salvio

Supercool Expansion (LO)

If supercooling is strong enough in a generic theory of the form \mathcal{L}^{ns} , to good accuracy, the full effective action for relevant values of χ can be described by **three and only three parameters**: χ_0 , $\bar{\beta}$ and g defined as

$$g^2 \chi^2 \equiv \sum_b n_b m_b^2(\chi) + \sum_f m_f^2(\chi).$$

The conditions for supercool expansion is verified if

$$\epsilon \equiv \frac{g^4}{6\bar{\beta} \log \frac{\chi_0}{T}} < 1.$$

- The LO of this expansion correspond to approximate the effective potential as

$$J_B(x) \approx J_B(0) + \frac{\pi^2}{12}x, \quad J_F(x) \approx J_F(0) - \frac{\pi^2}{24}x,$$

so

$$\bar{V}_{\text{eff}}(\chi, T) \approx \frac{m^2(T)}{2}\chi^2 - \frac{\lambda(T)}{4}\chi^4, \quad S_3 \approx c_3 \frac{m}{\lambda},$$

where

$$m^2(T) \equiv \frac{g^2 T^2}{12}, \quad \lambda(T) \equiv \bar{\beta} \log \frac{\chi_0}{T}, \quad c_3 = 18.8973 \dots$$

- It is possible to calculate several FOTP parameters:
 - ▶ **Nucleation temperature:** Defined as the temperature T_n for which

$$\Gamma(T_n) \approx H(T_n)^4 \equiv H_n^4.$$

- ▶ **Inverse duration:** Defined as

$$\beta \equiv \left[\frac{1}{\Gamma} \frac{d\Gamma}{dt} \right]_{t_n}.$$

Supercool Expansion

- The nucleation temperature T_n is obtained solving the equation

$$\Gamma(T_n) \approx H_n^4 \approx H_I^4 \quad \longrightarrow \quad cX - 4X^2 - a \approx 0$$

where $H_I = \frac{\sqrt{\bar{\beta}}\chi_0^2}{4\sqrt{3}\bar{M}_P}$ is the Hubble rate when the vacuum is still dominant respect the true vacuum, and

$$X \equiv \log \frac{\chi_0}{T_n}, \quad a \equiv \frac{c_3 g}{\sqrt{12\bar{\beta}}}, \quad c \equiv \log \frac{4\sqrt{3}\bar{M}_P}{\sqrt{\bar{\beta}}\chi_0} + \frac{3}{2} \log \frac{a}{2\pi}.$$

- Solving the equation we obtain

$$T_n \approx \chi_0 \exp \left(\frac{\sqrt{c^2 - 16a - c}}{8} \right).$$

- The inverse duration can be written as

$$\begin{aligned}\frac{\beta}{H_n} &\approx \left[T \frac{d}{dT} (S_3/T) - 4 - \frac{3}{2} T \frac{d}{dT} \log(S_3/T) \right]_{T=T_n} \\ &\approx \frac{a}{\log^2(\chi_0/T_n)} - 4 - \frac{3}{2} \frac{1}{\log(\chi_0/T_n)},\end{aligned}$$

neglecting the last term

$$\frac{\beta}{H_n} \approx \frac{a}{\log^2(\chi_0/T_n)} - 4.$$

- LO approximation means neglect terms of relative order $\sqrt{\epsilon}$
- The NLO expansion includes higher order terms in the expansion of the thermal function

$$J_B(x) \approx J_B(0) + \frac{\pi^2}{12}x - \frac{\pi}{6}x^{3/2}, \quad J_F(x) \approx J_F(0) - \frac{\pi^2}{24}x.$$

- The NLO effective potential includes a perturbative cubic term

$$\bar{V}_{\text{eff}}(\chi, T) \approx \frac{m^2(\chi)}{2}\chi^2 - \frac{k(T)}{3}\chi^3 - \frac{\lambda(T)}{4}\chi^4$$

$$k(T) \equiv \frac{\tilde{g}^3 T}{4\pi}, \quad \tilde{g}^3 \chi^3 \equiv \sum_b n_b m_b^3(\chi), \quad \tilde{g} \leq g.$$

Extending the validity of the supercool expansion

- The supercool expansion is still valid for $\epsilon \sim 1$ only if the number of degrees of freedom N with dominant coupling to the field χ is large

$$g \sim \sqrt{N}\tau, \quad \tilde{g} \lesssim \sqrt[3]{N}\tau \quad \longrightarrow \quad \tilde{g}^3/g^3 \lesssim 1/\sqrt{N}$$

- The cubic term in the effective potential gets suppressed by a factor $\lesssim 1/\sqrt{N}$
- $1/X = 6\bar{\beta}/\epsilon g^4$ is still small for $\epsilon \sim 1$
- Truncating the small- x expansion of the thermal function up to order $x^{3/2}$ because the higher-order terms involve smaller and smaller coefficients

Improved Supercool Expansion

- In general, when $\epsilon \sim 1$, in the effective potential **the cubic term cannot be treated as a perturbation** (i.e. the term of order $x^{3/2}$ is not perturbative in the expansion of the thermal functions)
- However we can **treat the cubic term as LO** and the other higher order term as a perturbation JCAP **12** (2023), 046 arXiv:2307.04694 [hep-ph], A. Salvio
- We can rewrite the effective potential and the 3d euclidean action at the LO of the **Improved Supercool Expansion** as

$$\tilde{V}_{\text{eff}}(\phi, T) = \frac{1}{2}\phi^2 - \frac{1}{3}\phi^3 - \frac{\tilde{\lambda}}{4}\phi^4, \quad S_3 = -\frac{8\pi m^3}{k^2} \int_0^\infty d\rho \rho^2 \tilde{V}_{\text{eff}}(\phi, T),$$

$$\tilde{\lambda}(T) \equiv \frac{\lambda m^2}{k^2} = \frac{(4\pi)^2 \bar{\beta}}{12 \tilde{g}^6 / g^2} \log(\chi_0/T) \geq \frac{2\pi^2}{9\epsilon}.$$

- We are interested for values of $\tilde{\lambda} \sim 1$: for such values we can write S_3 as
JHEP 02 (2023), 125, arXiv:2212.08085 [hep-ph], N. Levi, T. Opferkuch, D. Redigolo

$$S_3 = \frac{27\pi m^3}{2k^2} \frac{1 + \exp(-1/\sqrt{\tilde{\lambda}})}{1 + \frac{9}{2}\tilde{\lambda}}.$$

- This expression reproduces the numerical calculation at the $\sim 1\%$ level for the values of $\tilde{\lambda}$ we are interested in
- The validity of this expression of S_3 has been established in a **model independent way within the improved supercool expansion**

Improved Supercool Expansion

- Solving the equation $\Gamma(T_n) \approx H_n^4$

$$a_1 - a_2 \tilde{\lambda} = F(\tilde{\lambda}) \equiv \frac{1 + \exp(-1/\sqrt{\tilde{\lambda}})}{2/9 + \tilde{\lambda}},$$

$$a_1 \equiv \frac{c c_3 k^3}{3\pi a \bar{\beta} m^2}, \quad a_2 \equiv \frac{4c_3 k^4}{3\pi a \bar{\beta}^2 m^4}.$$

- We are interested in the smallest real and positive solution $\tilde{\lambda}_n(T) \equiv \tilde{\lambda}(T_n)$ for which the straight line $a_1 - a_2 \tilde{\lambda}$ reaches $F(\tilde{\lambda})$ from below in increasing $\tilde{\lambda}$ (if it exist)
- Following the same steps as before we can calculate β/H_n

$$\frac{\beta}{H_n} \approx \frac{\pi^3 g^5}{6\sqrt{3}\tilde{g}^8} \frac{(4\pi)^2 \bar{\beta}}{\tilde{g}^4} (-F'(\tilde{\lambda}_n)) - 4.$$

1 FOPT, Supercool Expansion and Improved Supercool Expansion

2 Late blooming mechanism

Late Blooming Mechanism

Small Overview

Phys.Rev.D 105 (2022) 2, L021303, J. Liu, L. Bian, R. G. Cai, Z. K. Guo and S. J. Wang

- Vacuum decay is a **probabilistic process**
- There can be some regions that **persist in the false vacuum** for a longer time than the background
- These regions can eventually **collapse into Primordial Black Holes (PBH)** if the density contrast reaches the critical value δ_c




Figure 3: Late Blooming Mechanism

Late Blooming Mechanism

Expanding the vacuum decay rate as

$$\Gamma(t) = \Gamma(t_n) \exp(\beta(t - t_n) + \beta_2(t - t_n)^2 + \dots) \approx H_n^4 e^{\beta(t - t_n)},$$

one can characterize the late blooming mechanism in supercooled 1OPT:

Phys.Rev.D 110 (2024) 4, 043514, arXiv:2305.04942 [hep-ph], Y. Gouttenoire, T. Volansky

Collapse Probability, PBH Fraction and PBH Mass

$$\mathcal{P}_{\text{coll}} \approx \exp \left[-a_{\mathcal{P}} \left(\frac{\beta}{H_n} \right)^{b_{\mathcal{P}}} (1 + \delta_c)^{c_{\mathcal{P}}} \frac{\beta}{H_n} \right], \quad \text{for } \alpha = \left(\frac{T_{\text{eq}}}{T_n} \right)^4 \gg 1, \quad \delta_c \simeq 0.45,$$

$$a_{\mathcal{P}} \approx 0.5646, \quad b_{\mathcal{P}} \approx 1.266, \quad c_{\mathcal{P}} \approx 0.6639,$$

$$f_{\text{PBH}} \approx \frac{\mathcal{P}_{\text{coll}}}{6.0 \times 10^{-12}} \frac{T_{\text{eq}}}{500 \text{GeV}}, \quad M_{\text{PBH}} \approx M_{\odot} \left(\frac{20}{g_*(T_{\text{eq}})} \right)^{1/2} \left(\frac{140 \text{MeV}}{T_{\text{eq}}} \right)^2.$$

It is also possible calculate the RMS Kerr Parameter of the PBHs
arXiv:2409.06494 [gr-qc], I. K. Banerjee, T. Harada

PBH Initial Spin

$$\sqrt{\langle a_*^2 \rangle} \approx \frac{2.1 \times 10^{-3}}{23.484 - 1.25 \log_{10}(f_{\text{PBH}}) - 1.25 \log_{10}\left(\frac{\Omega_{\text{CDM}}}{0.26}\right) - 0.625 \log_{10}\left(\frac{M_{\text{PBH}}}{10^{15} \text{g}}\right)}.$$

Late Blooming & Supercool Expansion

We can express the physical parameters of the Late Blooming Mechanism in terms of $(\chi_0, \bar{\beta}, g, \tilde{g})$

Late Blooming & Supercool Expansion

$$T_{\text{eq}}^4 \approx \frac{15\bar{\beta}\chi_0^4}{8\pi^2 g_*(T_{\text{eq}})}, \quad M_{\text{PBH}} \approx M_{\odot} \left(\frac{2\pi^2}{3\bar{\beta}} \right)^{1/2} \left(\frac{280\text{MeV}}{\chi_0} \right)^2, \quad H_n \sim H_I \approx \frac{\sqrt{\bar{\beta}}\chi_0^2}{4\sqrt{3}\bar{M}_P},$$

$$\frac{\beta}{H_n} \approx \frac{a}{\log^2(\chi_0/T_n)} - 4 \quad \text{(Non-Improved LO)},$$

$$\frac{\beta}{H_n} \approx \frac{\pi^3 g^5}{6\sqrt{3}\tilde{g}^8} \frac{(4\pi)^2 \bar{\beta}}{\tilde{g}^4} (-F'(\tilde{\lambda}_n)) - 4 \quad \text{(Improved LO)}.$$

Then we can calculate:

$$f_{\text{PBH}}(\beta/H_n(\chi_0, \bar{\beta}, g, \tilde{g})), \quad \sqrt{\langle a_*^2 \rangle} (f_{\text{PBH}}(\beta/H_n(\chi_0, \bar{\beta}, g, \tilde{g})), M_{\text{PBH}}(T_{\text{eq}}(\chi_0, \bar{\beta}))).$$

Late Blooming & Supercool Expansion

The expansion

$$\Gamma(t) = \Gamma(t_n) \exp(\beta(t - t_n) + \beta_2(t - t_n)^2 + \dots) \approx H_n^4 e^{\beta(t - t_n)},$$

can be justified on the ground of the supercool expansion. Using:

$$dt = -dT/(TH) \approx -dT/(TH_I) \longrightarrow T(t) \approx T_n e^{-H_I(t - t_n)},$$

for the supercool expansion at LO

$$\frac{S_3}{T} \approx c_3 \frac{m}{T\lambda} = \frac{c_3 g}{\sqrt{12} \bar{\beta} \log \frac{\chi_0}{T}} \equiv \frac{a}{X + \log \frac{T_n}{T}} \approx \frac{a}{X + H_I(t - t_n)},$$

$$X \equiv \log \frac{\chi_0}{T_n},$$

Late Blooming & Supercool Expansion

we get

$$\Gamma(t) \approx T^4(t) \exp(-S_3/T(t)) \approx T_n^4 \exp\left(-\frac{a}{X + H_I(t - t_n)} - 4H_I(t - t_n)\right).$$

Expanding for t around t_n

$$\frac{1}{1 + \frac{H_I(t - t_n)}{X}} = 1 - \frac{H_I(t - t_n)}{X} + \left(\frac{H_I(t - t_n)}{X}\right)^2 + \dots + (-1)^k \left(\frac{H_I(t - t_n)}{X}\right)^k + \dots$$

noting that $H_I = H_{\text{eq}}/\sqrt{2} = \gamma/(\sqrt{3}t_{\text{eq}})$

$$\frac{H_I(t - t_n)}{X} = \frac{\gamma(\tau - \tau_n)}{\sqrt{3}X}, \quad \gamma = 0.76329\dots, \quad \tau \equiv \frac{t}{t_{\text{eq}}},$$

Late Blooming & Supercool Expansion

- As long as $\tau - \tau_n$ is small respect to $\sqrt{3}X/\gamma$, $\Gamma(t) \approx H_n^4 e^{\beta(t-t_n)}$ is a **good approximation**.
- This holds also for the improved supercool expansion ($\epsilon \sim 1$ at $T = T_n$)

$$\bar{\beta} \sim \frac{g^4}{(4\pi)^2} \quad (\text{loop suppressed}) \quad \longrightarrow \quad X \sim 26 \quad \longrightarrow \quad \frac{\gamma}{\sqrt{3}X} \sim 10^{-2}$$

Late Blooming & Supercool Expansion




Figure 4: Improved Supercool at LO

Late Blooming & Supercool Expansion




Figure 5: Supercool at LO

Late Blooming & Supercool Expansion




Figure 6: Improved Supercool at LO

Late Blooming & Supercool Expansion




Figure 7: Supercool at LO

Late Blooming & Supercool Expansion




Figure 8: Improved LO




Figure 9: Improved LO

Late Blooming & Supercool Expansion




Figure 10: Improved Supercool LO




Figure 11: Improved Supercool LO

Late Blooming & Supercool Expansion




Figure 12: Supercool LO




Figure 13: Supercool LO

An example: B-L Model

- $U(1)_{B-L}$ model phenomenology can be described with Improved Supercool Expansion JCAP **12** (2023), 046 arXiv:2307.04694 [hep-ph], A. Salvio

$$\begin{aligned}\mathcal{L} = & \mathcal{L}_{\text{SM}}^{\text{ns}} + D_\mu A^\dagger D^\mu A + \bar{N}_j i \gamma_\mu D^\mu N_j - \frac{1}{4} B'_{\mu\nu} B'^{\mu\nu} \\ & + \left(Y_{ij} L_i \mathcal{H} N_j + \frac{1}{2} y_{ij} A N_i N_j + \text{h.c.} \right) \\ & - \lambda_a |A|^4 + \lambda_{ah} |A|^2 |\mathcal{H}|^2\end{aligned}$$

- The gauge group is $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ and

$$D_\mu = \partial_\mu + ig_3 T^\alpha G_\mu^\alpha + ig_2 T^a W_\mu^a + ig_Y \mathcal{Y} B_\mu + i[g_m \mathcal{Y} + g'_1 (B - L)] B'_\mu$$

An example: B-L Model

- the one-loop RG equation for the quartic coupling λ_a

$$(4\pi)^2 \mu \frac{d}{d\mu} \lambda_a = 96g_1'^4 - 48\lambda_a g_1'^2 + 20\lambda_a^2 + 2\lambda_{ah}^2 + 2\lambda_a \text{Tr}(yy^\dagger) - \text{Tr}(yy^\dagger yy^\dagger)$$

- Neglecting λ_{ah} and y (right-handed neutrino Majorana masses are taken below EW scale)

$$\bar{\beta} = \frac{96g_1'^4}{(4\pi)^2}$$

- The background dependent Z' mass is

$$M_{Z'}(\chi) = 2|g_1'|\chi$$

An example: B-L Model

- The collective couplings are:

$$g = 2\sqrt{3}|g'_1|, \quad \tilde{g} = 2\sqrt[3]{3}|g'_1| = \frac{g}{\sqrt[6]{3}},$$

we get

$$\bar{\beta} = \frac{2g^4}{3(4\pi)^2}$$

and

$$m_\chi = \sqrt{\frac{2}{3}} \frac{g^2}{4\pi} \chi_0, \quad M_h = \sqrt{\lambda_{ah}} \chi_0.$$

An example: B-L Model




Figure 14: Improved Supercool at LO

- The (improved) supercool expansion is a **powerful tool** to study the **phenomenology of FOPT**
- The FOPT phenomenology related to a **general RSB model** can be described by using **just few parameters** $(\chi_0, \bar{\beta}, g, \tilde{g})$
- We described using (improved) supercool expansion the production of PBH via late blooming mechanism, and provided a model that can account for an appreciable fraction of PBH




Figure 15: Energy Densities Late blooming




Figure 16: Density Contrast




Figure 17: Hubble rate Late Blooming Mechanism




Figure 18: Hubble Radius late blooming mechanism